APPLICATION OF NANOPARTICLES BASED ON HYDROPHILIC POLYMERS AS PHARMACEUTICAL FORMS

WO9804244 (A1)		Also published as:
1998-02-05		EP0860166 (A1)
PILAR [ES]; REMUNAN LOPES CARMEN [ES]; VILA JATO		EP0860166 (B1) PT860166 (T)
UNIV SANTIAGO COMPOSTELA [ES]; ALONSO FERNANDEZ MARIA JOSE [ES]; CALVO SALVE PILAR [ES]; REMUNAN LOPES CARMEN [ES]; VILA JATO JOSE LUIS [ES]		
		more >>
A61K9/51 ; A61K9/51 ; (IPC1-7): A61K9/51		Cited documents:
A61K9/51	(S)A	1105040700 (4)
WO1996ES00186 19961022 ES19960001685 19960729		US5346703 (A) WO9605810 (A1) WO9620698 (A2)
	1998-02-05 ALONSO FERNANDEZ MARIA JOSE [ES]; CALVO SALVE PILAR [ES]; REMUNAN LOPES CARMEN [ES]; VILA JATO JOSE LUIS [ES] UNIV SANTIAGO COMPOSTELA [ES]; ALONSO FERNANDEZ, MARIA JOSE [ES]; CALVO SALVE PILAR [ES]; REMUNAN LOPES CARMEN [ES]; VILA JATO JOSE LUIS [ES] **A61K9/51; A61K9/51; (IPC1-7): A61K9/51 A61K9/51 WO1996ES00186 19961022	1998-02-05 ALONSO FERNANDEZ MARIA JOSE [ES]; CALVO SALVE PILAR [ES]; REMUNAN LOPES CARMEN [ES]; VILA JATO JOSE LUIS [ES] UNIV SANTIAGO COMPOSTELA [ES]; ALONSO FERNANDEZ MARIA JOSE [ES]; CALVO SALVE PILAR [ES]; REMUNAN LOPES CARMEN [ES]; VILA JATO JOSE LUIS [ES] A61K9/51; A61K9/51; (IPC1-7): A61K9/51 A61K9/51 WO1996ES00186 19961022

Abstract of WO 9804244 (A1)

Application of nanoparticles based on hydrophilic polymers as, pharmaceutical forms for the administration of active macromolecules. The nanoparticles (having a nanometric size and a hydrophilic character), also called nanospheres or latex, are colloidal systems comprised of the combination of hydrophilic polymers and an active ingredient having a high molecular weight (active macromolecule, molecular weight higher than 1000 daltons). The hydrophilic polymers are the chitosane (an aminopolysaccharide) or its derivatives and the polyoxyethylene or its derivatives. The association of the active macromolecule to said nanoparticles takes place in an aqueous phase without having to use organic solvents or auxiliary toxic substances. The active ingredient charge capacity of the nanoparticles is extremely high and additionally said charge is released in a controlled and time extended way.; Additionally, said nanoparticles have a positive surface electric charge which the intensity may vary in relation to its composition.

Data supplied from the **esp@cenet** database — Worldwide